

Distinct Roles of P2X Receptors in Modulating Glutamate Release at Different Primary Sensory Synapses in Rat Spinal Cord

Terumasa Nakatsuka, Kenzo Tsuzuki, Jennifer X. Ling, Hideki Sonobe, and Jianguo G. Gu

McKnight Brain Institute of the University of Florida, Department of Oral Surgery Division of Neuroscience, College of Dentistry, University of Florida, Gainesville, Florida, 32610

Submitted 26 December 2002; accepted in final form 10 February 2003

Nakatsuka, Terumasa, Kenzo Tsuzuki, Jennifer X. Ling, Hideki Sonobe, and Jianguo G. Gu. Distinct roles of P2X receptors in modulating glutamate release at different primary sensory synapses in rat spinal cord. *J Neurophysiol* 89: 3243–3252, 2003; 10.1152/jn.01172.2002. Using spinal cord slice preparations and patch-clamp recordings in lamina II and lamina V regions, we tested a hypothesis that P2X receptor subtypes differentially modulate glutamate release from primary afferent terminals innervating different sensory regions. We found that activation of P2X receptors by α,β -methylene-ATP increased glutamate release onto >80% of DH neurons in both lamina regions. However, two distinct types of modulation, a transient and a long-lasting enhancement of glutamate release were observed. In lamina II recordings, >70% of the modulation was transient. In contrast, P2X receptor-mediated modulation was always long-lasting in lamina V. Pharmacologically, both transient and long-lasting types of modulation were blocked by 10 μ M pyridal-phosphate-6-azophenyl-2',4'-disulphonic acid tetrasodium, a broad-spectrum P2X receptor antagonist. Transient modulation was not observed in the presence of 1 μ M trinitrophenyl-ATP (TNP-ATP), a subtype-selective P2X receptor antagonist, suggesting that homomeric P2X₃ receptors may be involved in the transient modulation in lamina II. The long-lasting modulation remained in the presence of 1 μ M TNP-ATP. Selective removal of P2X₃-expressing afferent terminals by the targeting toxin saporin-conjugated isolectin B4 or surgical removal of superficial DH did not affect P2X receptor-mediated long-lasting modulation in lamina V. Taken together, these results suggest that P2X receptor subtypes play distinct roles in sensory processing in functionally different sensory regions.

INTRODUCTION

ATP P2X receptors (P2X) belong to a family of nonselective cation channels gated by extracellular ATP (Jahr and Jessell 1983; Krishtal et al. 1983; North and Surprenant 2000). P2X receptors can mediate fast excitatory synaptic transmission (Bardoni et al. 1997; Edwards et al. 1992; Evans et al. 1992). They also play roles in modulating glutamatergic, GABAergic, and glycinergic synaptic transmission in the CNS (Deuchars et al. 2001; Gu and MacDermott 1997; Hugel and Schlichter 2000; Jang et al. 2001; Khakh and Henderson 1998; Li et al. 1998; Rhee et al. 2000). The roles of P2X receptors in sensory signaling in the periphery have been established (Bland-Ward and Humphrey 1997; Cockayne et al. 2000; Hamilton et al. 1999; Souslova et al. 2000; Stanfa et al. 2000; Tsuda et al.

1999a,b, 2000). P2X activation also facilitates glutamate release from primary afferent central terminals in the spinal cord (Nakatsuka and Gu 2001), which implicates a central role of P2X and ATP in sensory processing.

Seven P2X subunits (P2X₁ to P2X₇) have been identified and cloned (North and Surprenant 2000). Six homomeric P2X subtypes (P2X₁ to P2X₅, P2X₇) and at least four heteromeric P2X subtypes (P2X₂₊₃, P2X₄₊₆, P2X₁₊₅, and P2X₂₊₆) can be formed by the seven subunits in heterologous expression systems (Khakh et al. 2001; also see Brown et al. 2002). These P2X subtypes can be classified by the ATP analogue, α,β -methylene-ATP ($\alpha\beta$ meATP), into $\alpha\beta$ meATP-sensitive and -insensitive subgroups. Homomeric P2X₁ and P2X₃ and heteromeric P2X₂₊₃, P2X₁₊₅ and P2X₄₊₆ are sensitive to $\alpha\beta$ meATP, and other subtypes have very low sensitivity to $\alpha\beta$ meATP (Khakh et al. 2001). $\alpha\beta$ meATP-sensitive P2X receptors can be further divided into rapidly desensitizing group, including homomeric P2X₁ and P2X₃ (Chen et al. 1995; Valera et al. 1994), and weakly desensitizing group including heteromeric P2X₂₊₃, P2X₁₊₅, and P2X₄₊₆ (Haines et al. 1999; Le et al. 1998; Lewis et al. 1995; Torres et al. 1998). Pyridal-phosphate-6-azophenyl-2',4'-disulphonic acid tetrasodium (PPADS) blocks most P2X receptors with little selectivity (Khakh et al. 2001). TNP-ATP selectively blocks P2X₁, P2X₃, and P2X₂₊₃ at nanomolar concentration range (Virginio et al. 1998).

Many dorsal root ganglion (DRG) neurons express P2X receptors (Jahr and Jessell 1983; Krishtal et al. 1983). Two types of currents, rapidly and weakly desensitizing currents, can be elicited by ATP or $\alpha\beta$ meATP (Grubb and Evans 1999; Li et al. 1999; Ueno et al. 1999). This indicates that at least two subtypes of functional P2X receptors are expressed on DRG neurons. Rapidly and weakly desensitizing P2X receptors may have distinct sensory functions in the periphery (Cook et al. 1997; Dowd et al. 1998; Sawynok and Reid 1997; Tsuda et al. 2000). Rapidly desensitizing currents are generally thought to be mediated by homomeric P2X₃; weakly desensitizing P2X currents are believed to be mainly mediated by P2X₂₊₃ (Burgard et al. 1999; Lewis et al. 1995; Tsuda et al. 2000; Xu and Huang 2002). However, other P2X subtypes may also mediate the weakly desensitizing currents in DRG neurons because

Address for reprint requests: J. G. Gu, McKnight Brain Institute of the University of Florida, Department of Oral Surgery Division of Neuroscience, College of Dentistry, University of Florida, Box 100416, Gainesville, Florida 32610 (E-mail: jgu@dental.ufl.edu).

The costs of publication of this article were defrayed in part by the payment of page charges. The article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

most P2X subunits are found to be expressed in sensory neurons (Collo et al. 1996; Xiang et al. 1998).

Spinal cord DH, the first central site for sensory processing, can be divided into structurally and functionally distinct lamina regions (Willis and Coggeshall 1991). Lamina II is a major receiving center for the inputs from fine afferent fibers that mainly carry nociceptive signals. In contrast, lamina V receives many convergent sensory inputs from both nociceptive and nonnociceptive afferent fibers (Willis and Coggeshall 1991). This region plays an important role in sensory hypersensitivity associated with allodynia and hyperalgesia. We have previously shown that P2X receptors are expressed at the central terminals of primary afferent fibers innervating lamina V neurons and that activation of these receptors modulates sensory transmission to lamina V neurons (Nakatsuka and Gu 2001). However, the properties of these P2X receptors on afferent central terminals have not been examined in detail. In lamina II, little is known about the involvement of P2X receptors in regulating glutamate release from the central terminal of primary afferent fibers. For these reasons, in the present study we have characterized P2X receptor-mediated modulation of glutamate release in both lamina II and lamina V. Further, the present study demonstrates that the modulation in these two regions is distinct due to the differential expression of P2X receptor subtypes at the central terminals of primary afferent fibers.

METHODS

Spinal-cord slice preparation

Principles of laboratory animal care (National Institutes of Health publication No. 86-23, revised 1985) were followed in all the experiments described in this study. Transverse spinal cord slices were prepared from L₅ spinal cords of rats at the postnatal age of 14–21 days as described previously (Nakatsuka and Gu 2001). Briefly, in each experiment, a rat was anesthetized with isoflurane, and then a laminectomy was performed. The lumbosacral spinal cord was taken out and placed in the preoxygenated Krebs solution at 2–4°C. After cutting all ventral and dorsal roots near the root entry zone, the pia-arachnoid membrane was removed, and the spinal cord was mounted on a Vibratome. Unless otherwise indicated, the spinal cord was cut into transverse slices each in the thickness of 500 μm . A spinal cord slice was transferred to a recording chamber (~0.5 ml) and placed on the stage of an upright IR-DIC microscope (BX50WI; Olympus, Tokyo). The slice was superfused with Krebs solution at flow rate of 10 ml/min at room temperature (22°C). The Krebs solution contained (in mM) 117 NaCl, 3.6 KCl, 2.5 CaCl₂, 1.2 MgCl₂, 1.2 NaH₂PO₄, 25 NaHCO₃, and 11 glucose; the solution was saturated with 95% O₂-5% CO₂ and had pH of 7.3.

Patch-clamp recordings from spinal cord slices

Lamina regions were identified under a $\times 10$ objective, and individual neurons were identified with a $\times 40$ objective under IR-DIC microscope (Olympus BX50). The microscope was coupled with a CCD camera (CCD100; DAGE-MTI, Michigan City, IN) and a video monitor screen. Whole cell patch-clamp recordings were made from DH neurons with microelectrodes filled with a solution containing (in mM) 135 K⁺-gluconate, 5 KCl, 0.5 CaCl₂, 2 MgCl₂, 5 EGTA, and 5 HEPES, pH 7.3; the electrode resistance was ~5 M Ω after filling the electrode solution. Signals were amplified with Axopatch 200B (Axon Instruments, Union City, CA), filtered at 2 kHz, sampled at 5 kHz using pCLAMP 7 (Axon Instruments), and stored in a personal computer. Stable recordings could be obtained from spinal cord slices

and maintained for several hours. Spontaneous excitatory postsynaptic currents (sEPSCs) and miniature EPSCs (mEPSCs) were recorded at holding potential of -60 mV where GABAergic and glycinergic synaptic currents were negligible (Nakatsuka and Gu 2001). In some experiments, s- and mEPSCs were recorded in the presence of bicuculline (20 μM) and strychnine (2 μM). sEPSCs were recorded in the absence of TTX and mEPSCs in the presence of 0.5 μM TTX. Effects of $\alpha\beta\text{meATP}$ on sEPSCs and mEPSCs were examined in lamina II and lamina V neurons. $\alpha\beta\text{meATP}$ was puff-applied as well as bath-applied in lamina II recordings; $\alpha\beta\text{meATP}$ was bath-applied in lamina V recordings. For puff application, pressure microinjection of 1 mM $\alpha\beta\text{meATP}$ was applied at 69 kPa for 5 s by a Picospitzer (General Valve, Fairfield, NJ). Puff electrodes were positioned with their tips being located at a distance of ~150 μm to the soma of the recorded neurons in lamina II. Puff pipettes were fabricated in the identical way as the recording micropipettes. All antagonists tested were applied through bath solution for ≥ 10 min before the applications of $\alpha\beta\text{meATP}$. Synaptic events including mEPSCs and sEPSCs were analyzed using Mini Analysis Program (Jaevin Software, Anderson Place, GA) with criteria being the same as previously described (Gu and MacDermott 1997). In analyzing the change of sEPSC and mEPSC frequency after bath application of $\alpha\beta\text{meATP}$, the time course of sEPSC and mEPSC frequency before and after $\alpha\beta\text{meATP}$ was first constructed with time bin of 10 s. Then the average response in three bins (30 s) around the peak was used to calculate the changes in reference to the control.

Chemical removal of IB4-positive sensory terminals

In one set of experiments, electrophysiological experiments were performed on the spinal cord slice preparations obtained from rats that were preinjected with saporin-conjugated isolectin B4 (IB4-saporin) to the sciatic nerves. IB4-saporin (Advanced Targeting System, San Diego, CA) injection was performed on rats 5–7 days prior to electrophysiological recordings and immunostaining. In brief, rats were deeply anesthetized with isoflurane, and the left sciatic nerve was isolated. IB4-saporin (2 μl , 1 mg/ml) was injected into the sciatic nerve via a microelectrode (~0.5 μm in diameter) under a dissecting microscope. The wound was washed with 5 ml saline and sutured with 4-0 monofilament nylons. Transverse spinal cord slices (250 μm in thickness) were prepared from L₅ spinal cords 5–7 days after the injection. Whole cell patch-clamp recordings were then made from DH neurons of the injected side. After the electrophysiological recordings, spinal cord slices were fixed, and immunostaining was performed on these spinal cord slices. In another group of experiment, recordings were conducted on lamina V neurons in the spinal cord slices whose superficial laminae were surgically removed.

Immunohistochemistry

IB4 staining and P2X₃ subunit immunostaining were performed to determine the effectiveness of IB4-saporin in removing IB4-positive/P2X₃-expressing sensory terminals in the spinal cord. VR1-ir was also examined in some slices after recordings. In brief, the transverse spinal cord slices (250 μm in thickness) were put in a 35-mm petri dish and fixed with 4% paraformaldehyde (PFA, in PBS buffer solution) for 12 h at 4°C after the electrophysiological recordings. The slices were transferred into a 4% PFA solution containing 0.4% Triton X-100 and incubated at 4°C for 3 h. They were washed three times with PBS and then mounted onto glass slides and allow to air-dry. They were then encircled with hydrophobic resin (PAP Pen). For IB4 staining, slices were incubated with IB4-biotin (Sigma, St. Louis, MO) at room temperature for 1 h. After a rinse with 1% goat serum PBS solution three times, the slices were further incubated with Streptavidin-Alexa 350 (Molecular Probes, Eugene, OR). For P2X₃ immunostaining, slices were incubated with a polyclonal guinea pig anti-P2X₃ receptor antibody (1:3,000; Neuromics, Minneapolis, MN)

over night at 4°C. After a rinse with 1% goat serum PBS solution three times, the slices were further incubated with a secondary antibody for 3 h at room temperature. The secondary antibody (1:100 in goat serum PBS solution) was a goat anti-guinea pig IgG-conjugated with Alexa-488 (Molecular Probes). The slices were washed three times with 1% goat serum PBS solution. For VR1 immunostaining, slices were incubated with a polyclonal rabbit anti-VR1 receptor antibody (1:3,000; Neuromics) overnight at 4°C. After a rinse with 1% goat serum PBS solution three times for 20 min, the slices were further incubated with a secondary antibody for 3 h at room temperature. The secondary antibody (1:100 in goat serum PBS solution) was a goat anti-rabbit IgG conjugated with Alexa-594 (Molecular Probes). Slices were washed three times with 1% goat serum PBS solution and then coverslip was applied with a glycerol-based anti-photobleach medium. Slices were viewed under an inverted fluorescent microscope (IX-70, Olympus, Tokyo, Japan) or a confocal microscope (1024 ES, Bio-Rad, Hercules, CA) equipped with a krypton-argon laser and T1 and T2A filter blocks, attached with an inverted fluorescent microscope (IX-70, Olympus). To provide a quantitative measure of the changes of P2X₃-ir in spinal cord slice sections after IB4-injection, the images were first converted to a gray scale for which the darker gray represents heavier immunostaining and the lighter gray represents less immunostaining. The immunostaining intensity was then quantitatively measured by the Scion-Imaging Program (Scion, Frederick, MD). The immunostaining intensity in lamina VII was taken as a reference of negative staining. The ratio between the immunostaining intensity in lamina III and lamina VII was calculated and used for the comparison of P2X₃-ir.

$\alpha\beta$ meATP, PPADS, capsaicin, bicuculline, and strychnine were purchased from Sigma. 6-cyano-7-nitroquinoxaline-2,3-dion (CNQX) and tetrodotoxin (TTX) were purchased from Tocris (Ballwin, MO). Trinitrophenyl-ATP (TNP-ATP) was purchased from Molecular Probe. Data represent means \pm SE. Paired Student's *t*-test were used for statistical comparison, and significance was considered at the *P* < 0.05 level.

RESULTS

P2X receptor-mediated modulation of glutamate release onto lamina II neurons

The metabolically stable ATP analogue $\alpha\beta$ meATP was used in this study because its action in spinal cord slice preparations was found to be mainly on primary afferent central terminals, and it had little effect on DH neurons (Nakatsuka and Gu 2001). Puff application (5 s) was used to rapidly deliver $\alpha\beta$ meATP ($\sim 0.1 \mu\text{M}$) to lamina II near the places where the recorded lamina II neurons were located. Although the concentration of $\alpha\beta$ meATP inside the puff electrode was 1 mM, its actual concentration that reached afferent central terminals should be much lower because of instant dilution by surrounding buffer solution. sEPSC frequency was increased in 9 of 11 lamina II neurons recorded (82%, Fig. 1, A, B, and E) after a 5-s puff application of $\alpha\beta$ meATP. The responses had very rapid onset. The effects were not due to mechanical force by puff pressure because puff application of bath solution did not produce any change in sEPSC frequency. The overall changes in sEPSC frequency were $359 \pm 119\%$ of controls ($n = 11$). Of the nine cells showing the increased sEPSC frequency, seven of them showed transient responses that were terminated 1–2 s before the end of 5-s $\alpha\beta$ meATP application (Fig. 1A). The remaining two neurons showed responses that lasted for >10 s with the 5-s $\alpha\beta$ meATP application (Fig. 1B). Similar to its effects on sEPSC frequency, mEPSC frequency in seven of the nine lamina II neurons recorded also was increased after puff




FIG. 1. Increases of spontaneous and miniature excitatory postsynaptic current (sEPSC and mEPSC) frequency in lamina II neurons following puff application of α,β -methylene-ATP ($\alpha\beta$ meATP). A: an example shows a transient increase of sEPSC frequency in a lamina II neuron after puff application of $\alpha\beta$ meATP. The response was desensitized before the end of a 5-s $\alpha\beta$ meATP application. $\alpha\beta$ meATP concentration inside puff electrode was 1 mM. The line above the trace indicates the duration of $\alpha\beta$ meATP application. B: an example shows a prolonged increase of sEPSC frequency in a different lamina II neuron after puff application of 1 mM $\alpha\beta$ meATP. The response was sustained during a 5-s $\alpha\beta$ meATP application and lasted for several seconds after the application. C and D: puff applications of $\alpha\beta$ meATP (1 mM) produced a transient increase (C) and a prolonged increase (D) of mEPSC frequency in the presence of 500 nM TTX in 2 lamina II neurons. TTX was continuously present in bath solution and was also included in puff solution. E and F: pooled results show $\alpha\beta$ meATP-induced increases of sEPSC (E) and mEPSC (F) frequency in lamina II neurons. sEPSC frequency was increased in 9 of 11 lamina II neurons recorded (E). Of these 9 cells, 7 cells had transient and 2 cells had the prolonged responses. mEPSC frequency was increased in 7 out of 9 lamina II cells recorded (F). Of these 7 cells, 5 cells had transient and 2 cells had the prolonged responses.

application of $\alpha\beta$ meATP (Fig. 1, C, D, and F). The overall changes in mEPSC frequency were $437 \pm 147\%$ ($n = 9$). Of the seven cells showing the increased mEPSC frequency, five of them showed transient responses that were terminated 1–2 s

before the end of 5-s $\alpha\beta$ meATP application (Fig. 1C). The remaining two neurons showed responses that lasted for >10 s with 5-s $\alpha\beta$ meATP application (Fig. 1D). In the preceding experiments, $\alpha\beta$ meATP did not directly evoke P2X receptor-mediated whole cell currents from the recorded DH neurons because there was little change in baseline current levels during $\alpha\beta$ meATP applications (Fig. 1, A–D). The sEPSCs and mEPSCs in the preceding experiments were mediated by the release of glutamate from terminals connecting to the recorded neurons because they were abolished completely in the presence of 10 μ M 6-cyano-7-nitroquinoxaline-2,3-dione (CNQX, not shown). It was noted that when a cell had a transient response to $\alpha\beta$ meATP, a second application of $\alpha\beta$ meATP in the same cell usually could not induce an increase in the frequency of sEPSCs and mEPSCs even when the time interval between two applications was >10 min. $\alpha\beta$ meATP-induced transient responses rarely recovered completely, even with very long time wash in normal bath solution (not shown). This phenomenon is consistent with the previous findings that some P2X receptors such as homomeric P2X₃ receptors recover very slowly from desensitization (Khakh et al. 2001).

The effects of two P2X receptor antagonists, PPADS and TNP-ATP, on $\alpha\beta$ meATP-induced responses were tested in lamina II recordings. Due to the very slow and incomplete recovery from the desensitization of $\alpha\beta$ meATP-induced responses in most recordings, $\alpha\beta$ meATP was applied only one time in the presence of an antagonist in each recording. Therefore the effects of antagonists were analyzed by group comparison. This approach is valid because $\sim 80\%$ of cells responded to $\alpha\beta$ meATP in the absence of P2X antagonists. In the presence of 10 μ M PPADS in bath perfusion solution, puff application of $\alpha\beta$ meATP (1 mM $\alpha\beta$ meATP mixed with 10 μ M PPADS inside puff electrode) did not significantly increase sEPSC frequency ($102 \pm 4\%$ of control) in any of nine neurons recorded in lamina II (Fig. 2, A and D). Similar experiments were performed to test the effects of TNP-ATP. In 10 neurons recorded in the presence of 1 μ M TNP-ATP, puff application of 1 mM $\alpha\beta$ meATP (1 mM $\alpha\beta$ meATP mixed with 1 μ M TNP-ATP inside puff electrode) did not significantly increase sEPSC frequency in 8 neurons (Fig. 2B). Two neurons clearly showed an increase of sEPSC frequency (Fig. 2C), but the responses in these two cells were different from the transient responses observed in most lamina II neurons. They had longer-lasting responses that continue for several seconds after the termination of the 5-s $\alpha\beta$ meATP application. The average response induced by $\alpha\beta$ meATP in the presence of 1 μ M TNP-ATP is presented in Fig. 2D. The average response includes data from eight nonresponsive and two responsive neurons.

We examined the modulation of glutamate release in lamina II after the prolonged P2X receptor activation. In this experiment, 100 μ M $\alpha\beta$ meATP was applied through bath solution for a period of 60 s. Of 22 cells recorded in lamina II, only 5 of them (23%) showed long-lasting increases of sEPSC frequency after $\alpha\beta$ meATP application (Fig. 3, A and C). The responses lasted for >150 s (Fig. 3A). In the remaining 17 cells (77%) recorded, neither transient nor long-lasting response was observed (Fig. 3, B and C). We further examined the effects of $\alpha\beta$ meATP on mEPSCs in the presence of 0.5 μ M TTX. Similar to the effects on sEPSC frequency, bath application of $\alpha\beta$ meATP only produced long-lasting increases of mEPSC




FIG. 2. Effects of pyridyl-phosphate-6-azophenyl-2',4'-disulphonic acid tetrasodium (PPADS) and trinitrophenyl-ATP (TNP-ATP) on $\alpha\beta$ meATP-induced changes of sEPSC frequency in lamina II. A: a sample trace shows the lack of response to a 5-s puff application of 1 mM $\alpha\beta$ meATP when 10 μ M PPADS was present. B: a sample trace shows the lack of response to a 5-s puff application of 1 mM $\alpha\beta$ meATP in a lamina II cell when 1 μ M TNP-ATP was present. C: in a different lamina II cell, a 5-s puff application of 1 mM $\alpha\beta$ meATP produced a prolonged increase of sEPSC frequency in the presence of 1 μ M TNP-ATP. D: pooled results show the overall responses to the puff applications of 1 mM $\alpha\beta$ meATP in normal bath ($\alpha\beta$ meATP, $n = 9$), in the presence of PPADS ($\alpha\beta$ meATP/PPADS, $n = 9$), and in the presence of TNP-ATP ($\alpha\beta$ meATP/TNP-ATP, $n = 10$). The pooled responses in the presence of TNP-ATP include 8 cells as illustrated in B and 2 cells as illustrated in C. Data represent means \pm SE; * $P < 0.05$, Student's *t*-test.

frequency in a small percent of neurons recorded (4 of 18 neurons, 22%, Fig. 3, D and F). The majority of lamina II neurons recorded (14 of 18 cells, 78%) showed no change in mEPSC frequency (Fig. 3, E and F).

P2X receptor-mediated modulation of glutamate release onto lamina V neurons

Although P2X receptor-mediated modulation of glutamate release onto lamina V neurons was reported previously (Nakatsuka and Gu 2001), the properties of this modulation have not been characterized in detail. The following experiments were designed to compare P2X receptor-mediated synaptic modulation in lamina V with that in lamina II.




FIG. 3. Changes in the frequency of sEPSCs and mEPSCs in lamina II neurons after the prolonged bath application of $\alpha\beta\text{meATP}$. *A*: a histogram shows the time course of the long-lasting increase of sEPSC frequency in a lamina II neuron after bath application of $100\ \mu\text{M}$ $\alpha\beta\text{meATP}$ for 60 s. The line above the histogram indicates the duration of $\alpha\beta\text{meATP}$ application. *B*: an example shows that bath application of $100\ \mu\text{M}$ $\alpha\beta\text{meATP}$ for 60 s did not produce any significant increase of sEPSC frequency in a lamina II neuron. *C*: pooled results from 22 lamina II cells. Five cells had long-lasting responses to $\alpha\beta\text{meATP}$ as illustrated in *A* and the remaining 17 cells had little response. Each line connecting 2 solid circles represents a recording from a lamina II neuron before and after the application of $\alpha\beta\text{meATP}$. *D–F*: the experiments were the same as that shown in *A–C* except mEPSCs were recorded in the presence of 500 nM TTX ($n = 18$). $\alpha\beta\text{meATP}$ -induced long-lasting increases of mEPSC frequency were only observed in 4 lamina II neurons (*F*). In all the recordings for sEPSCs and mEPSCs, no transient response was observed after bath applications of $\alpha\beta\text{meATP}$.

Different from most recordings in lamina II, >90% of lamina V neurons showed long-lasting increases of sEPSC and mEPSC frequency (Fig. 4) after bath application of $100\ \mu\text{M}$ $\alpha\beta\text{meATP}$ for 60 s. Of 22 cells recorded for sEPSCs (Fig. 4, *A* and *B*), 20 cells showed the increased sEPSC frequency by $\alpha\beta\text{meATP}$. The effects lasted for $210 \pm 14\ \text{s}$ ($n = 20$), about three times longer than the period of $\alpha\beta\text{meATP}$ application. The remaining two cells did not show a change in sEPSC frequency. The overall changes of sEPSC frequency were $371 \pm 48\%$ of control ($n = 22$) after $\alpha\beta\text{meATP}$ application. The effects of $100\ \mu\text{M}$ $\alpha\beta\text{meATP}$ on mEPSCs were tested in 20 cells (Fig. 4, *C* and *D*), 19 of them (95%) showed the increased mEPSC frequency. The effects lasted for $181 \pm 15\ \text{s}$. Only one cell showed no response. The overall changes in mEPSC frequency were $342 \pm 41\%$ of control ($n = 20$) after bath application of $100\ \mu\text{M}$ $\alpha\beta\text{meATP}$. The increases of sEPSC and mEPSC frequency could be reproduced by multiple applications of $100\ \mu\text{M}$ $\alpha\beta\text{meATP}$ at a time interval of 10 min (not shown, but see Fig. 5*B*). $\alpha\beta\text{meATP}$ at the concentration of $10\ \mu\text{M}$ was tested in eight lamina V neurons; these neurons were also tested with $100\ \mu\text{M}$ $\alpha\beta\text{meATP}$ (Fig. 4*E*). mEPSC frequency was increased to $168 \pm 27\%$ of control ($P < 0.05$, $n = 8$) with $10\ \mu\text{M}$ $\alpha\beta\text{meATP}$ and to $376 \pm 54\%$ of control with $100\ \mu\text{M}$ $\alpha\beta\text{meATP}$ ($P < 0.05$, $n = 8$).

The two P2X receptor antagonists, PPADS and TNP-ATP,

were examined to determine antagonist profiles of $\alpha\beta\text{meATP}$ -induced responses in lamina V neurons. Because $\alpha\beta\text{meATP}$ -induced increases of mEPSCs and sEPSCs showed essentially identical responses (Fig. 4), TTX was omitted from bath solution, and sEPSCs were measured in all of the following experiments. The $\alpha\beta\text{meATP}$ -induced increases of sEPSC frequency were sensitive to the block by PPADS. As shown in Fig. 5, *A* and *C*, the increases of sEPSC frequency by $100\ \mu\text{M}$ $\alpha\beta\text{meATP}$ ($403 \pm 89\%$ of control, $n = 6$) were completely abolished in the presence of $10\ \mu\text{M}$ PPADS ($104 \pm 5\%$ of control, $n = 6$). In contrast to PPADS, TNP-ATP did not significantly inhibit $\alpha\beta\text{meATP}$ -induced increases of sEPSCs (Fig. 5*B,C*). Of six lamina V neurons recorded, $\alpha\beta\text{meATP}$ ($100\ \mu\text{M}$) facilitated sEPSC frequency to $472 \pm 89\%$ of control in the absence of TNP-ATP; $\alpha\beta\text{meATP}$ still increased sEPSC frequency to $448 \pm 99\%$ of control in the same neurons when $1\ \mu\text{M}$ TNP-ATP was present (Fig. 5, *B* and *C*).

alpha-beta-meATP-induced long-lasting increases of glutamate release onto lamina V neurons after the removal of P2X3-expressing afferent terminals

P2X₃-ir in the spinal cord section was restricted in inner lamina II (Fig. 6*B*), and overlapped with IB4-positive primary afferent central terminals (Fig. 6*A*). However, many lamina V




FIG. 4. Changes in the frequency of sEPSCs and mEPSCs in lamina V neurons after the prolonged bath application of $\alpha\beta$ meATP. *A*: a histogram shows a long-lasting increase of sEPSC frequency in a lamina V neuron after bath application of $100\ \mu\text{M}$ $\alpha\beta$ meATP for 60 s. *B*: pooled results show that the long-lasting increases of sEPSC frequency occurred in almost all neurons recorded in lamina V ($n = 22$). *C* and *D*: similar to *A* and *B* except the experiments were conducted in the presence of 500 nM TTX. $\alpha\beta$ meATP induced long-lasting increases of mEPSC frequency in almost all lamina V neurons recorded ($n = 20$). *E*: $\alpha\beta$ meATP-induced increases of mEPSC frequency at the concentrations of $10\ \mu\text{M}$ and $100\ \mu\text{M}$ of $\alpha\beta$ meATP in lamina V neurons ($n = 8$).

neurons are known to extend their dendrites dorsally into superficial laminae, making it possible that P2X_{2+3} -expressing afferent terminals were involved (North 2002). To address whether or not P2X_{2+3} receptors were responsible for the long-lasting modulation in lamina V recordings, we examined effects of $\alpha\beta$ meATP on sEPSC frequency after chemically destroying P2X_3 -expression afferent neurons using IB4-saporin (Fig. 6). Five days after the injection of $2\ \mu\text{l}$ IB4-saporin ($1\ \text{mg/ml}$) into the sciatic nerve, IB4 staining in the ipsilateral side disappeared (Fig. 6A) from the spinal cord slice preparations of L_5 section. We used the ratio of P2X_3 -ir intensity between lamina II and lamina VII as a quantitative measure for the change of P2X_3 -ir after IB4-saporin injection. The ratio of P2X_3 -ir intensity was 3.00 ± 0.43 ($n = 6$) in the control sides, and it became 1.01 ± 0.11 ($n = 6$, $P < 0.05$) in the IB4-injected side. Thus P2X_3 -ir was almost completely lost in the ipsilateral side (Fig. 6B).

Electrophysiological experiments were performed in such slices prior to immunostaining of P2X_3 receptors. The removal of P2X_3 -expression terminals in the recorded slice sections was confirmed in each experiment after recordings (Fig. 6B). $\alpha\beta$ meATP-induced responses in lamina V on the ipsilateral site showed similar responses to those recordings from slices of un-injected rats. As shown in Fig. 6, *C* and *D*, bath application of $100\ \mu\text{M}$ $\alpha\beta$ meATP still largely enhanced sEPSC frequency to $305 \pm 42\%$ of control ($n = 20$). The effects were observed in almost all lamina V neurons (90%, 18 of 20 neurons from 6 slices) in the IB4-saporin injected sides (Figs. 6, *C* and *D*, and *7D*).

To further confirm that a subtype of P2X receptors other than P2X_{2+3} receptors was responsible for the long-lasting modulation in lamina V recordings, we examined $\alpha\beta$ meATP-induced responses in lamina V neurons after the removal of superficial DH (lamina I–III; Fig. 7). A second purpose of this




FIG. 5. Effects of PPADS and TNP-ATP on $\alpha\beta\text{meATP}$ -induced changes of sEPSC frequency in lamina V. *A*: a histogram shows the changes of sEPSC frequency in a lamina V neuron after 60-s bath applications of 100 μM $\alpha\beta\text{meATP}$ in the absence and presence of 10 μM PPADS. *B*: a histogram shows the changes of sEPSC frequency in a lamina V neuron after 60-s bath applications of 100 μM $\alpha\beta\text{meATP}$ in the absence and presence of 1 μM TNP-ATP. *C*: pooled results show the overall responses to bath applications of 100 μM $\alpha\beta\text{meATP}$ in the absence of antagonists ($\alpha\beta\text{meATP}$, $n = 6$), in the presence of 10 μM PPADS ($\alpha\beta\text{meATP}/\text{PPADS}$, $n = 6$), or in the presence of 1 μM TNP-ATP ($\alpha\beta\text{meATP}/\text{TNP-ATP}$, $n = 6$). Control was sEPSC frequency before $\alpha\beta\text{meATP}$ application and was scaled as 100%. Data represent means \pm SE; * $P < 0.05$, paired Student's *t*-test.

experiment was to determine the location (superficial vs. deep laminae) of these $\alpha\beta\text{meATP}$ -sensitive synapses. Figure 7, *A* and *B*, illustrates the position of P2X₃-ir and its relative position to the portion of the dorsal horn that was surgically removed. In addition, Fig. 7*A* also showed that P2X₃-ir largely overlapped with VR1-ir in the inner lamina II (Fig. 7*A*). The removal of superficial lamina abolished capsaicin-induced increases of sEPSC frequency in lamina V neurons ($n = 10$, not

shown) (also see Nakatsuka et al. 2002), indicating the effectiveness of removing VR1/P2X₃-expressing terminals. When $\alpha\beta\text{meATP}$ (100 μM) was tested in such preparation, bath




FIG. 6. $\alpha\beta\text{meATP}$ -induced long-lasting increases of sEPSC frequency in lamina V neurons after the removal of P2X₃-expressing afferent terminals by saporin-conjugated isolectin B4 (IB4-saporin). *A*: the micrograph shows IB4 staining in lamina II of control side (right side) and the lack of IB4 staining in the left side in a L₅ spinal cord slice section. The slice was obtained from a rat for which IB4-saporin (IB4-SAP) was injected into the left sciatic nerve. The L₅ spinal cord slice section was obtained and IB4 staining performed 5 days after the injection of IB4-SAP. *B*: the experiment was the same as *A* except immunostaining was performed for P2X₃ subunit. P2X₃-ir was shown in lamina II of right side (control side) and disappeared from left side (IB4-SAP injected side). *C*: an example shows a recording from a lamina V neuron in the IB4-SAP injected side. $\alpha\beta\text{meATP}$ still induced long-lasting increases of sEPSC frequency in the lamina V neuron when P2X₃-expressing terminals were removed by IB4-SAP. *D*: pooled results from 22 lamina V neurons of IB4-SAP injected sides. The removal of IB4/P2X₃-expressing terminals had little effect on $\alpha\beta\text{meATP}$ -induced responses in lamina V. In all experiments, the removal of P2X₃-expressing terminals in the IB4-SAP-injected sides was confirmed by P2X₃ immunostaining after each electrophysiological experiment.




FIG. 7. $\alpha\beta\text{meATP}$ -induced long-lasting increases of sEPSC frequency in lamina V after surgical removal of superficial laminae. *A*: a confocal image of immunoreactivity for P2X₃ subunit and VR1 receptor in a spinal cord slice. P2X₃-ir (green) was restricted in lamina II and VR1-ir (red) across both lamina I and lamina II regions. In lamina II, P2X₃-ir and VR1-ir overlapped (yellow) substantially. *B*: the micrograph shows a spinal cord slice whose superficial laminae on the right side were removed surgically. A recording electrode is shown in the field, and the tip of the electrode is located in lamina V region on the right side. An inset in *B* is a resized image of *A* to show the relative position of P2X₃-ir to the surgically removed portion of the dorsal horn. *C*: $\alpha\beta\text{meATP}$ -induced increases of sEPSC frequency in the preparation as illustrated in *B* ($n = 12$). *D*: a summary shows $\alpha\beta\text{meATP}$ -induced changes of sEPSC frequency in intact spinal cord slice ($n = 22$), in the slices whose superficial laminae were surgically removed ($n = 12$), and in the slice obtained from IB4-SAP injected rats ($n = 20$). Data represent means \pm SE; Student's *t*-test were used for comparison, and no significant differences were found between the intact slice group and 2 other groups.

application of $\alpha\beta\text{meATP}$ still largely increased sEPSC frequency in all 12 lamina V neurons recorded (Fig. 7, *C* and *D*). The overall changes of sEPSC frequency were $320 \pm 89\%$ of control ($n = 12$), similar to the responses in intact slices. Thus the removal of P2X₃-expressing terminals either chemically or surgically did not significantly change the $\alpha\beta\text{meATP}$ -induced responses in lamina V neurons (Fig. 7*D*).

DISCUSSION

We have demonstrated that P2X receptors play distinct roles in modulating glutamate release at different primary sensory synapses in the dorsal horn of the spinal cord. A transient type of modulation, mediated by rapidly desensitizing subtype of P2X receptors, is predominant in the primary sensory synapses of lamina II. Whereas in lamina V, a long-lasting type of modulation is engaged in almost all the primary sensory synapses and is mediated by weakly desensitizing subtype of P2X receptors. These results implicate that P2X receptor subtypes play distinct roles in central processing of different primary

sensory inputs. Because both lamina II and V are involved in nociceptive transmission, the facilitation of glutamate release by presynaptic P2X receptors may have implications in the changes of synaptic plasticity during physiological and pathological pain states.

P2X receptor-mediated modulation of glutamate release onto lamina II neurons

Immunochemical evidence has shown the expression of P2X subunits at the central terminals of primary afferent fibers (Guo et al. 1999; Vulchanova et al. 1996–1998). Although activation of P2X receptors produced glutamate release from sensory synapses in a DGR-DH co-culture system (Gu and MacDermott 1997; Labrakakis et al. 2000), only one study suggests that P2X receptors may play a role in modulating glutamate release in lamina II (Li et al. 1998). Using puff application to rapidly deliver $\alpha\beta\text{meATP}$, we found that the frequency of sEPSCs and mEPSCs, i.e., synaptic release of glutamate, was increased in most recordings in lamina II. Most responses were shown to be transient and rapidly desensitized. In contrast, most recordings did not reveal any significant increase in the frequency of sEPSCs and mEPSCs when $\alpha\beta\text{meATP}$ was bath applied. The lack of transient responses during bath application of $\alpha\beta\text{meATP}$ to the tissue, which inactivated the rapidly desensitizing P2X receptors before it could be activated. Thus most afferent terminals connecting to lamina II neurons expressed rapidly desensitizing P2X receptors, and activation of these receptors produces a transient increase of glutamate release. Pharmacologically, we showed that the transient responses in lamina II were sensitive to the block by both PPADS and TNP-ATP. Because P2X₃ subunits are high expressed at the afferent terminals innervating lamina II regions (Vulchanova et al. 1997, 1998) and homomeric P2X₃ receptors are highly sensitive to the block by TNP-ATP (Virginio et al. 1998; also see Burgard et al. 1999; Grubb and Evans 1999), the transient responses induced by $\alpha\beta\text{meATP}$ in lamina II were most likely mediated by homomeric P2X₃ receptors. However, the potential contribution of P2X₁ receptors cannot be excluded.

We have shown that in a small population of lamina II neurons, focal and bath application of $\alpha\beta\text{meATP}$ induced a prolonged increase of glutamate release. This suggests that the weakly desensitizing P2X receptors are expressed on a small number of afferent terminals innervating lamina II. In DRG neurons, weakly desensitizing P2X receptors that could be activated by $\alpha\beta\text{meATP}$ were highly speculated to be P2X₂₊₃ receptors (Burgard et al. 1999; Lewis et al. 1995; Xu and Huang 2002). P2X₂₊₃ receptors are sensitive to the block by TNP-ATP in nanomolar concentration range (Virginio et al. 1998). However, we found that during continuous perfusion of 1 μM TNP-ATP, $\alpha\beta\text{meATP}$ still could produce a long-lasting increase of glutamate release in a small number of lamina II neurons. This raises a possibility that some afferent central terminals innervating lamina II regions may express a P2X subtype different from P2X₂₊₃. It should be noted that our experimental protocol with TNP-ATP in this study could not allow us to determine the potential role of P2X₂₊₃ heteromeric receptors (Xu and Huang 2002) in $\alpha\beta\text{meATP}$ -induced long-lasting responses because TNP-ATP were present through the experiments in lamina II recordings. P2X₂₊₃-like currents have

recently been demonstrated in the soma of DRG neurons (Burgard et al. 1999; Grubb and Evans 1999; Rae et al. 1998; Tsuda et al. 2000; Ueno et al. 1999) although co-expression of P2X₂ and P2X₃ subunits at the same central terminals of primary afferent fibers was not clearly evident immunocytochemically with confocal microscopic study (Vulchanova et al. 1997).

P2X receptor-mediated modulation of glutamate release onto lamina V neurons

$\alpha\beta$ meATP largely increased sEPSC and mEPSC frequency in almost all lamina V neurons for a prolonged period. This effect was distinct from $\alpha\beta$ meATP-induced transient responses recorded in lamina II cells. These results suggest that weakly desensitizing P2X receptors are expressed at the central terminals of primary afferents innervating lamina V neurons. $\alpha\beta$ meATP-induced increases of mEPSCs and sEPSCs in lamina V were unlikely due to its action on DH neurons because $\alpha\beta$ meATP-sensitive P2X receptors were found on DH inhibitory interneurons not on glutamatergic DH neurons (Hugel and Schlichter 2000; Jang et al. 2001; Rhee et al. 2000). The expression of $\alpha\beta$ meATP-sensitive P2X receptors on A δ -afferent terminals in lamina V has been more directly demonstrated recently by the effects of $\alpha\beta$ meATP on A δ -afferent-evoked EPSCs recorded in lamina V neurons; $\alpha\beta$ meATP at low concentrations potentiated A δ -afferent-evoked EPSCs (Nakatsuka and Gu 2001) and failed the evoked EPSCs in lamina V neurons at high concentrations (Tsuzuki et al. 2003).

It has been widely thought that P2X receptors on the central terminals of A δ -afferent fibers are P2X₂₊₃ receptors (North 2002). However, $\alpha\beta$ meATP-induced responses in lamina V recordings were remained in the presence of 1 μ M TNP-ATP, raising a possibility that these weakly desensitizing P2X receptors were not P2X₂₊₃ receptors. One possibility for the lack of TNP-ATP effects could be due to its metabolic instability in tissue preparations (Lewis et al. 1998). However, the abolishment of transient responses in lamina II suggested that TNP-ATP reached targets at a concentration high enough to completely block homomeric P2X₃ receptors. Because P2X₂₊₃ receptors were shown to be slightly less sensitive to TNP-ATP antagonism than P2X₃ receptors (Virginio et al. 1998), we used two TNP-ATP-independent approaches to test the idea that those $\alpha\beta$ meATP-activated P2X receptors were not P2X₂₊₃. One approach was to chemically remove P2X₃-expressing afferent fibers, and the other approach was to surgically remove superficial laminae. P2X₃ subunits have been shown to be predominantly, if not exclusively, expressed on IB4-positive DRG neurons (Vulchanova et al. 1998). IB4-positive DRG neurons could be selectively destroyed by IB4-saporin (IB4-SAP) (Vulchanova et al. 2001). We found that IB4-staining was negative and P2X₃-ir was lost in L₅ spinal slice sections after sciatic nerve injections of IB4-SAP. In such a spinal cord slice preparation, we found that $\alpha\beta$ meATP still produced long-lasting increases of glutamate release onto lamina V neurons. Furthermore, $\alpha\beta$ meATP-induced responses were not altered in lamina V recordings when P2X₃-ir regions (superficial laminae) in the spinal cord were surgically removed. The results with surgical removal of superficial laminae also indicate that afferent terminals expressing the weakly desensitizing P2X receptors make synaptic contacts with lamina V neurons in

deep laminae. Taken together, the P2X receptors that mediated long-lasting enhancement of glutamate release onto lamina V neurons were unlikely to be P2X₂₊₃ receptors. The pharmacological profiles of $\alpha\beta$ meATP-mediated synaptic modulation in lamina V appear to be consistent with the involvement of P2X₁₊₅ and/or P2X₄₊₆. We have recently shown the similar pharmacological profiles of $\alpha\beta$ meATP-evoked currents directly recorded from dissociated DRG neurons (see Tsuzuki et al. 2003). However, in view that unidentified P2X heteromeric receptors (North 2002) may be present in native cells, it remains to be possible that P2X receptors on A δ -afferent terminals innervating lamina V region are new P2X receptor subtypes.

In conclusion, differential modulation of sensory inputs into different sensory regions by P2X receptor subtypes represents an important mechanism of sensory processing in the spinal cord dorsal horn.

This work was supported by a National Institute of Neurological Disorders and Stroke Grant NS-38254 to J. G. Gu.

REFERENCES

- Bardoni R, Goldstein PA, Lee CJ, Gu JG, and MacDermott AB. ATP P2X receptors mediate fast synaptic transmission in the dorsal horn of the rat spinal cord. *J Neurosci* 17: 5297–2304, 1997.
- Bland-Ward PA and Humphrey PP. Acute nociception mediated by hind-paw P2X receptor activation in the rat. *Br J Pharmacol* 122: 365–371, 1997.
- Brown SG, Townsend-Nicholson A, Jacobson KA, Burnstock G, and King BF. Heteromultimeric P2X(1/2) receptors show a novel sensitivity to extracellular pH. *J Pharmacol Exp Ther*. 300: 673–680, 2002.
- Burgard EC, Niforatos W, van Biesen T, Lynch KJ, Touma E, Metzger RE, Kowaluk EA, and Jarvis MF. P2X receptor-mediated ionic currents in dorsal root ganglion neurons. *J Neurophysiol* 82: 1590–1598, 1999.
- Chen CC, Akopian AN, Sivilotti L, Colquhoun D, Burnstock G, and Wood JN. A P2X purinoceptor expressed by a subset of sensory neurons. *Nature* 377: 428–431, 1995.
- Cockayne DA, Hamilton SG, Zhu QM, Dunn PM, Zhong Y, Novakovic S, Malmberg AB, Cain G, Berson A, Kassotakis L, Hedley L, Lachnit WG, Burnstock G, McMahon SB, and Ford AP. Urinary bladder hyporeflexia and reduced pain-related behavior in P2X3-deficient mice. *Nature* 407: 1011–1015, 2000.
- Collo G, North RA, Kawashima E, Merlo-Pich E, Neidhart S, Surprenant A, and Buell G. Cloning OF P2X5 and P2X6 receptors and the distribution and properties of an extended family of ATP-gated ion channels. *J Neurosci* 16: 2495–2507, 1996.
- Cook SP, Vulchanova L, Hargreaves KM, Elde R, and McCleskey EW. Distinct ATP receptors on pain-sensing and stretch-sensing neurons. *Nature* 387: 505–508, 1997.
- Deuchars SA, Atkinson L, Brooke RE, Musa H, Milligan CJ, Batten TF, Buckley NJ, Parson SH, and Deuchars J. Neuronal P2X7 receptors are targeted to presynaptic terminals in the central and peripheral nervous systems. *J Neurosci* 21: 7143–7152, 2001.
- Dowd E, McQueen DS, Chessell IP, and Humphrey PP. P2X receptor-mediated excitation of nociceptive afferents in the normal and arthritic rat knee joint. *Br J Pharmacol* 125: 341–346, 1998.
- Edwards FA, Gibb AJ, and Colquhoun D. ATP receptor-mediated synaptic currents in the central nervous system. *Nature* 359: 144–147, 1992.
- Evans RJ, Derkach V, and Surprenant A. ATP mediates fast synaptic transmission in mammalian neurons. *Nature* 357: 503–505, 1992.
- Grubb BD and Evans RJ. Characterization of cultured dorsal root ganglion neuron P2X receptors. *Eur J Neurosci* 11: 149–154, 1999.
- Gu JG and MacDermott AB. Activation of ATP P2X receptors elicits glutamate release from sensory neuron synapses. *Nature* 389: 749–753, 1997.
- Guo A, Vulchanova L, Wang J, Li X, and Elde R. Immunocytochemical localization of the vanilloid receptor 1 (VR1): relationship to neuropeptides, the P2X3 purinoceptor and IB4 binding sites. *Eur J Neurosci* 11: 946–958, 1999.

- Haines WR, Torres GE, Voigt MM, and Egan TM.** Properties of the novel ATP-gated ionotropic receptor composed of the P2X(1) and P2X(5) isoforms. *Mol Pharmacol* 56: 720–727, 1999.
- Hamilton SG, Wade A, and McMahon SB.** The effects of inflammation and inflammatory mediators on nociceptive behaviour induced by ATP analogues in the rat. *Br J Pharmacol* 126: 326–332, 1999.
- Hugel S and Schlichter R.** Presynaptic P2X receptors facilitate inhibitory GABAergic transmission between cultured rat spinal cord dorsal horn neurons. *J Neurosci* 20: 2121–2130, 2000.
- Jahr CE and Jessell TM.** ATP excites a subpopulation of rat dorsal horn neurons. *Nature* 304: 730–733, 1983.
- Jang IS, Rhee JS, Kubota H, Akaike N, and Akaike N.** Developmental changes in P2X purinoceptors on glycinergic presynaptic nerve terminals projecting to rat substantia gelatinosa neurons. *J Physiol* 536: 505–519, 2001.
- Khakh BS and Henderson G.** ATP receptor-mediated enhancement of fast excitatory neurotransmitter release in the brain. *Mol Pharmacol* 54: 372–378, 1998.
- Khakh BS, Burnstock G, Kennedy C, King BF, North RA, Seguela P, Voigt M, and Humphrey PP.** International union of pharmacology. XXIV. Current status of the nomenclature and properties of P2X receptors and their subunits. *Pharmacol Rev* 53: 107–118, 2001.
- Krishtal OA, Marchenko SM, and Pidoplichko VI.** Receptor for ATP in the membrane of mammalian sensory neurons. *Neurosci Lett* 35: 41–45, 1983.
- Labrakakis C, Gerstner E, and MacDermott AB.** Adenosine triphosphate-evoked currents in cultured dorsal root ganglion neurons obtained from rat embryos: desensitization kinetics and modulation of glutamate release. *Neuroscience* 101: 1117–1126, 2000.
- Le KT, Villeneuve P, Ramjaun AR, McPherson PS, Beaudet A, and Seguela P.** Sensory presynaptic and widespread somatodendritic immunolocalization of central ionotropic P2X ATP receptors. *Neuroscience* 83: 177–190, 1998.
- Lewis C, Neidhart S, Holy C, North RA, Buell G, and Surprenant A.** Coexpression of P2X2 and P2X3 receptor subunits can account for ATP-gated currents in sensory neurons. *Nature* 377: 432–435, 1995.
- Lewis CJ, Surprenant A, and Evans RJ.** 2', 3'-O-(2, 4, 6-trinitrophenyl) adenosine 5'-triphosphate (TNP-ATP)—a nanomolar affinity antagonist at rat mesenteric artery P2X receptor ion channels. *Br J Pharmacol* 124: 1463–1466, 1998.
- Li C, Peoples RW, Lanthorn TH, Li ZW, and Weight FF.** Distinct ATP-activated currents in different types of neurons dissociated from rat dorsal root ganglion. *Neurosci Lett* 263: 57–60, 1999.
- Li P, Calejesan AA, and Zhuo M.** ATP P2x receptors and sensory synaptic transmission between primary afferent fibers and spinal dorsal horn neurons in rats. *J Neurophysiol* 80: 3356–3360, 1998.
- Nakatsuka T, Furue H, Yoshimura M, and Gu JG.** Activation of central terminal VR1 receptors and α meATP-sensitive P2X receptors reveals a converged synaptic activity onto the deep dorsal horn neurons of the spinal cord. *J Neurosci* 22: 1228–1237, 2002.
- Nakatsuka T and Gu JG.** ATP P2X receptor-mediated enhancement of glutamate release and evoked EPSCs in dorsal horn neurons of the rat spinal cord. *J Neurosci* 21: 6522–6531, 2001.
- North RA.** Molecular physiology of P2X receptors. *Physiol Rev* 82: 1013–1067, 2002.
- North RA and Surprenant A.** Pharmacology of cloned P2X receptors. *Annu Rev Pharmacol Toxicol* 40: 563–80, 2000.
- Petruska JC, Cooper BY, Johnson RD, and Gu JG.** Distribution patterns of different P2x receptor phenotypes in acutely dissociated dorsal root ganglion neurons of adult rats. *Exp Brain Res* 134: 126–132, 2000.
- Rhee JS, Wang ZM, Nabekura J, Inoue K, and Akaike N.** ATP facilitates spontaneous glycinergic IPSC frequency at dissociated rat dorsal horn interneuron synapses. *J Physiol* 524: 471–483, 2000.
- Rae MG, Rowan EG, and Kennedy C.** Pharmacological properties of P2X3-receptors present in neurons of the rat dorsal root ganglia. *Br J Pharmacol* 124: 176–180, 1998.
- Sawynok J and Reid A.** Peripheral adenosine 5'-triphosphate enhances nociception in the formalin test via activation of a purinergic P2X receptor. *Eur J Pharmacol* 330: 115–121, 1997.
- Stanfa LC, Kontinen VK, and Dickenson AH.** Effects of spinally administered P2X receptor agonists and antagonists on the responses of dorsal horn neurones recorded in normal, carrageenan-inflamed and neuropathic rats. *Br J Pharmacol* 129: 351–359, 2000.
- Souslova V, Cesare P, Ding Y, Akopian AN, Stanfa L, Suzuki R, Carpenter K, Dickenson A, Boyce S, Hill R, Nebunius-Oosthuizen D, Smith AJ, Kidd EJ, and Wood JN.** Warm-coding deficits and aberrant inflammatory pain in mice lacking P2X3 receptors. *Nature* 407: 1015–1017, 2000.
- Torres GE, Haines WR, Egan TM, and Voigt MM.** Co-expression of P2X₁ and P2X₅ receptor subunits reveals a novel ATP-gated ion channel. *Mol Pharmacol* 54: 989–993, 1998.
- Tsuda M, Koizumi S, Kita A, Shigemoto Y, Ueno S, and Inoue K.** Mechanical allodynia caused by intraplantar injection of P2X receptor agonist in rats: involvement of heteromeric P2X2/3 receptor signaling in capsaicin-insensitive primary afferent neurons. *J Neurosci* 20: RC90, 2000.
- Tsuda M, Ueno S, and Inoue K.** In vivo pathway of thermal hyperalgesia by intrathecal administration of alpha, beta-methylene ATP in mouse spinal cord: involvement of the glutamate-NMDA receptor system. *Br J Pharmacol* 127: 449–456, 1999a.
- Tsuda M, Ueno S, and Inoue K.** Evidence for the involvement of spinal endogenous ATP and P2X receptors in nociceptive responses caused by formalin and capsaicin in mice. *Br J Pharmacol* 128: 1497–1504, 1999b.
- Tsuzuki K, Ase A, Séguela P, Nakatsuka T, Wang C-Y, She J-X, and Gu JG.** TNP-ATP-resistant P2X ionic current on the central terminals and somata of rat primary sensory neurons. *J Neurophysiol* 89: 3235–3242, 2003.
- Ueno S, Tsuda M, Iwanaga T, and Inoue K.** Cell type-specific ATP-activated responses in rat dorsal root ganglion neurons. *Br J Pharmacol* 126: 429–436, 1999.
- Valera S, Hussy N, Evans RJ, Adams N, North RA, Surprenant A, and Buell G.** A new class of ligand-gated ion channel defined by P2x receptor for extracellular ATP. *Nature* 371: 516–519, 1994.
- Virginio C, Robertson G, Surprenant A, and North RA.** Trinitrophenyl-substituted nucleotides are potent antagonists selective for P2X1, P2X3, and heteromeric P2X2/3 receptors. *Mol Pharmacol* 53: 969–973, 1998.
- Vulchanova L, Arvidsson U, Riedl M, Wang J, Buell G, Surprenant A, North RA, and Elde R.** Differential distribution of two ATP-gated channels (P2X receptors) determined by immunocytochemistry. *Proc Natl Acad Sci USA* 93: 8063–8067, 1996.
- Vulchanova L, Olson TH, Stone LS, Riedl MS, Elde R, and Honda CN.** Cytotoxic targeting of isolectin IB4-binding sensory neurons. *Neuroscience* 108: 143–155, 2001.
- Vulchanova L, Riedl MS, Shuster SJ, Buell G, Surprenant A, North RA, and Elde R.** Immunohistochemical study of the P2X2 and P2X3 receptor subunits in rat and monkey sensory neurons and their central terminals. *Neuropharmacology* 36: 1229–1242, 1997.
- Vulchanova L, Riedl MS, Shuster SJ, Stone LS, Hargreaves KM, Buell G, Surprenant A, North RA, and Elde R.** P2X3 is expressed by DRG neurons that terminate in inner lamina II. *Eur J Neurosci* 10: 3470–3478, 1998.
- Willis WD Jr and Coggeshall RE.** *Sensory Mechanisms of the Spinal Cord*. New York: Plenum, 1991, p. 79–151.
- Xiang Z, Bo X, and Burnstock G.** Localization of ATP-gated P2X receptor immunoreactivity in rat sensory and sympathetic ganglia. *Neurosci Lett* 256: 105–108, 1998.
- Xu GY and Huang LY.** Peripheral inflammation sensitizes P2X receptor-mediated responses in rat dorsal root ganglion neurons. *J Neurosci* 22: 93–102, 2002.